

Inverse Functions

Consider function $f(x)$:




Inverse Functions

Consider function $f(x)$:



Can we reverse the process?




Inverse Functions

Consider function $f(x)$:



Can we reverse the process?



Sometimes yes, sometimes no.

Inverse Functions



Consider two functions $f(x) = 2x$ and $g(x) = \frac{x}{2}$

- they are *inverse functions*

Inverse Functions

Consider two functions $f(x) = 2x$ and $g(x) = \frac{x}{2}$




- they are *inverse functions*



Inverse Functions

Consider two functions $f(x) = 2x$ and $g(x) = \frac{x}{2}$




- they are *inverse functions*



Inverse Functions

Consider two functions $f(x) = 2x$ and $g(x) = \frac{x}{2}$

- they are *inverse functions*



Inverse Functions

Consider two functions $f(x) = 2x$ and $g(x) = \frac{x}{2}$

Let's check their compositions:

$$(f \circ g)(x)$$

and

$$(g \circ f)(x)$$

Inverse Functions

Consider two functions $f(x) = 2x$ and $g(x) = \frac{x}{2}$

Let's check their compositions:

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{x}{2}\right) = 2 \cdot \frac{x}{2} = x$$

and

$$(g \circ f)(x) = g(f(x)) = g(2x) = \frac{2x}{2} = x$$

Inverse Functions

[Def] Let f and g be the two functions such that

$$f(g(x)) = x \text{ for every } x \in D_g$$

and

$$g(f(x)) = x \text{ for every } x \in D_f.$$

The function g is the inverse of the function f and is denoted by f^{-1} (“ f -inverse”).

Inverse Functions

[Def] Let f and g be the two functions such that

$$f(g(x)) = x \text{ for every } x \in D_g$$

and

$$g(f(x)) = x \text{ for every } x \in D_f.$$

The function g is the inverse of the function f and is denoted by f^{-1} (“ f -inverse”).

Thus $f(f^{-1}(x)) = x$ and $f^{-1}(f(x)) = x$

The domain $D_{f^{-1}} = \text{range of } f$ and vice versa.

Inverse Functions

Example: let's verify that each function is inverse of the other:

$$f(x) = 3x + 8$$

$$g(x) = \frac{x - 8}{3}$$

Inverse Functions

Example: let's verify that each function is inverse of the other:

$$f(x) = 3x + 8 \qquad g(x) = \frac{x - 8}{3}$$

We need to check that $f(g(x)) = x$ and $g(f(x)) = x$:

Inverse Functions

Example: let's verify that each function is inverse of the other:

$$f(x) = 3x + 8 \qquad g(x) = \frac{x - 8}{3}$$

We need to check that $f(g(x)) = x$ and $g(f(x)) = x$:

$$f(g(x)) = f\left(\frac{x - 8}{3}\right) = 3 \left(\frac{x - 8}{3}\right) + 8 = x - 8 + 8 = x$$

Inverse Functions

Example: let's verify that each function is inverse of the other:

$$f(x) = 3x + 8$$

$$g(x) = \frac{x - 8}{3}$$

We need to check that $f(g(x)) = x$ and $g(f(x)) = x$:

$$f(g(x)) = f\left(\frac{x - 8}{3}\right) = 3 \frac{x - 8}{3} + 8 = x - 8 + 8 = x$$

$$g(f(x)) = g(3x + 8) = \frac{(3x + 8) - 8}{3} = \frac{3x}{3} = x$$

Inverse Functions

Example: let's verify that each function is inverse of the other:

$$f(x) = 3x + 8 \qquad g(x) = \frac{x - 8}{3}$$

We need to check that $f(g(x)) = x$ and $g(f(x)) = x$:

$$f(g(x)) = f\left(\frac{x - 8}{3}\right) = 3 \frac{x - 8}{3} + 8 = x - 8 + 8 = x \quad \checkmark$$

$$g(f(x)) = g(3x + 8) = \frac{(3x + 8) - 8}{3} = \frac{3x}{3} = x \quad \checkmark$$

Answer: yes, they are inverses of each other

In-class practice

Exercise 1:

Determine if $f(x)$ and $g(x)$ are inverse functions.

$$f(x) = \frac{1}{x} - 7$$

$$g(x) = \frac{1}{x+7}$$

In-class practice

Exercise 2:

Determine if $f(x)$ and $g(x)$ are inverse functions.

$$f(x) = x - 7 \qquad g(x) = 7x$$

Inverse Functions

Functions given by tables

Sometimes functions are given in a tabular format to us.

x	-2	0	1	4	7
$f(x)$	28	49	-9	15	6

Inverse Functions

Functions given by tables

Sometimes functions are given in a tabular format to us.

x	-2	0	1	4	7
$f(x)$	28	49	-9	15	6

We can find $f^{-1}(28), f^{-1}(49), f^{-1}(-9), f^{-1}(6), f^{-1}(15)$ using the table.

Inverse Functions

Functions given by tables

Sometimes functions are given in a tabular format to us.

x	-2	0	1	4	7
$f(x)$	28	49	-9	15	6

We can find $f^{-1}(28), f^{-1}(49), f^{-1}(-9), f^{-1}(6), f^{-1}(15)$ using the table.

We can invert $f(x)$ using the table

Inverse Functions

Functions given by tables

Sometimes functions are given in a tabular format to us.

x	-2	0	1	4	7
$f(x)$	28	49	-9	15	6

We can find $f^{-1}(28), f^{-1}(49), f^{-1}(-9), f^{-1}(6), f^{-1}(15)$ using the table.

We can invert $f(x)$ using the table

x	-9	6	15	28	49
$f^{-1}(x)$					

Inverse Functions

Functions given by tables

Sometimes functions are given in a tabular format to us.

x	-2	0	1	4	7
$f(x)$	28	49	-9	15	6

We can find $f^{-1}(28), f^{-1}(49), f^{-1}(-9), f^{-1}(6), f^{-1}(15)$ using the table.

We can invert $f(x)$ using the table

x	-9	6	15	28	49
$f^{-1}(x)$	1	7	4	-2	0

Inverse Functions

Finding the inverse of a function

If we are given an equation of a function $f(x)$, then we can find the inverse of $f(x)$ following these steps:

- 1) in the equation for $f(x)$, replace $f(x)$ with y
- 2) interchange x and y
- 3) solve for y

* if the equation doesn't define y as a function of x , then function f doesn't have an inverse function

* otherwise, the resulting equation defines an inverse function f^{-1}

- 4) if f has inverse function, replace y in step 3) by $f^{-1}(x)$
- 5) check: we can verify that $f(f^{-1}(x)) = x$ and $f^{-1}(f(x)) = x$

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

replace $f(x)$ with y

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

$$y = 5x - 9 \quad \text{replace } f(x) \text{ with } y$$

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

$$y = 5x - 9$$

replace $f(x)$ with y
interchange x and y

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

$$y = 5x - 9 \quad \text{replace } f(x) \text{ with } y$$

$$x = 5y - 9 \quad \text{interchange } x \text{ and } y$$

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

$$y = 5x - 9 \quad \text{replace } f(x) \text{ with } y$$

$$x = 5y - 9 \quad \text{interchange } x \text{ and } y$$

$$x + 9 = 5y \quad \text{solve for } y$$

$$\frac{x+9}{5} = y$$

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

$$y = 5x - 9 \quad \text{replace } f(x) \text{ with } y$$

$$x = 5y - 9 \quad \text{interchange } x \text{ and } y$$

$$x + 9 = 5y \quad \text{solve for } y$$

$$\frac{x+9}{5} = y \quad \text{defines a function of } x$$

Inverse Functions

Example: let's find an inverse of $f(x) = 5x - 9$

$$f(x) = 5x - 9$$

$$y = 5x - 9 \quad \text{replace } f(x) \text{ with } y$$

$$x = 5y - 9 \quad \text{interchange } x \text{ and } y$$

$$x + 9 = 5y \quad \text{solve for } y$$

$$\frac{x + 9}{5} = y \quad \text{defines a function of } x$$

$$f^{-1}(x) = \frac{x + 9}{5} \quad \text{replace } y \text{ with } f^{-1}(x)$$

In-class practice

Exercise 1: find an inverse, if it exists, of $f(x) = \sqrt[3]{2x+3}$

In-class practice

Exercise 1: find an inverse, if it exists, of $f(x) = \sqrt[3]{2x+3}$

replace $f(x)$ with y
interchange x and y
solve for y
defines a function of x ?
replace y with $f^{-1}(x)$

In-class practice

Exercise 2: find an inverse, if it exists, of $f(x) = 3x^2 - 10$

replace $f(x)$ with y
interchange x and y
solve for y
defines a function of x ?
replace y with $f^{-1}(x)$

Inverse Functions




The horizontal line test and one-to-one functions

A function f , has an inverse function f^{-1} if there is no horizontal line that intersects the graph of f at more than one point.

Inverse Functions

The horizontal line test and one-to-one functions




A function f , has an inverse function f^{-1} if there is no horizontal line that intersects the graph of f at more than one point.



Inverse Functions

The horizontal line test and one-to-one functions

A function f , has an inverse function f^{-1} if there is no horizontal line that intersects the graph of f at more than one point.



Inverse Functions

The horizontal line test and one-to-one functions


A function f , has an inverse function f^{-1} if there is no horizontal line that intersects the graph of f at more than one point.

[Def] A *one-to-one function* is a function in which no two different ordered pairs have the same second component, i.e. if $x_1 \neq x_2$ then $f(x_1) \neq f(x_2)$

Inverse Functions

Graphs of f and f^{-1}

Consider the graphs of $f(x) = x^3$ and $g(x) = \sqrt[3]{x}$




the graphs of inverse functions are reflections of each other about the line $y = x$.

Inverse Functions

Graphs of f and f^{-1}

Consider the graphs of $f(x) = x^3$ and $g(x) = \sqrt[3]{x}$



The coordinates of points in graphs of inverse functions are switched: see the point (1.26, 2) on the graph of x^3 and the point (2, 1.26) on the graph of $\sqrt[3]{x}$.

x	$y = x^3$
0	0
1	1
2	8
3	27

x	$y = \sqrt[3]{x}$
0	0
1	1
8	2
27	3

Homework assignment

- 1) zyBooks:** *review Section 1.7 Inverse Functions*
or
Textbook: *review Section 1.8 Inverse Functions.*
- 2)** We will have **Quiz 5** based on today's topics in the beginning of our next meeting.
- 3) WeBWorK:** **HW 5** (due date is in one week)